

**BLUE BIRD AUREO.5/BOV ROG  
TYPE B Pasture Cattle Feed  
Medicated**

For increased rate of weight gain and for control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline in pasture cattle over 700 pounds (slaughter, stocker, feeder cattle, beef replacement heifers).

**Active Drug Ingredients\***

Chlortetracycline ..... 1100 to 80000 g/ton  
Lasalocid ..... 600 to 80000 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Mixing Directions**

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 25 to 1100 grams chlortetracycline per ton and 30 to 600 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Body Wt. lb.	Feed Consumption lbs/head/day	Type B Concentration g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
		CTC	Lasalocid			CTC	Lasalocid
700	1	14000	12000	100	1900	700	600
	3.5	2000	1143	200	1800	200	114.3
	14	1000	600	100	1900	50	30
1000	2	5000	1500	200	1800	500	150
	4	5000	2000	100	1900	250	100

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.