

**BLUE BIRD AUREO/BOV FE  
TYPE B Confined Cattle Feed  
Medicated**

For improved feed efficiency and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle fed in confinement for slaughter.

**Active Drug Ingredients\***

Chlortetracycline ..... 4000 to 80000 g/ton  
Lasalocid ..... 100 to 80000 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Mixing Directions**

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 500 to 2000 grams chlortetracycline per ton and 10 to 30 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Body Wt. lb.	Feed Consumption lbs/head/day	Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
		CTC	Lasalocid			CTC	Lasalocid
500	10	10000	200	200	1800	1000	20
	10	20000	600	100	1900	1000	30
	15	6670	140	200	1800	667	14
	15	13340	600	100	1900	667	30
1000	20	10000	100	200	1800	1000	10
	20	20000	600	100	1900	1000	30
	30	6670	100	200	1800	667	10
	30	13340	480	100	1900	667	24

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.

**BLUE BIRD AUREO/BOV FE/ROG  
TYPE B Confined Cattle Feed  
Medicated**

For improved feed efficiency, increased rate of weight gain and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle fed in confinement for slaughter.

**Active Drug Ingredients\***

Chlortetracycline ..... 4000 to 80000 g/ton  
Lasalocid ..... 100 to 80000 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Mixing Directions**

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 500 to 1200 grams chlortetracycline per ton and 25 to 30 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Body Wt. lb.	Feed Consumption lbs/head/day	Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
		CTC	Lasalocid			CTC	Lasalocid
800	20	8000	250	200	1800	800	25
	20	16000	600	100	1900	800	30
	24	6670	250	200	1800	667	25
	24	13340	600	100	1900	667	30
1000	20	10000	250	200	1800	1000	25
	20	20000	600	100	1900	1000	30
	25	8000	250	200	1800	800	25
	25	16000	560	100	1900	800	28

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.

**BLUE BIRD AUREO/BOV ROG  
TYPE B Pasture Cattle Feed  
Medicated**

For increased rate of weight gain and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in pasture cattle (slaughter, stocker, feeder cattle, dairy and beef replacement heifers).

**Active Drug Ingredients\***

Chlortetracycline .....4000 to 80000 g/ton  
Lasalocid .....400 to 80000 g/ton

**Guaranteed Analysis**

Crude protein, not less than .....%  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than .....%  
Crude fiber, not more than .....%  
Calcium, not less than .....%  
Calcium, not more than .....%  
Phosphorus, not less than .....%  
Potassium, not less than .....%  
Salt<sup>1</sup>, not less than .....%  
Salt<sup>1</sup>, not more than .....%  
Sodium<sup>2</sup>, not less than .....%  
Sodium<sup>2</sup>, not more than .....%  
Vitamin A<sup>1</sup>, not less than .....IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Mixing Directions**

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 500 to 4000 grams chlortetracycline per ton and 30 to 600 grams lasalocid per ton. The following table provides examples of mixing rates and amounts\*\*:

Body Wt. lb.	Feed Consumption lbs/head/day	Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
		CTC	Lasalocid			CTC	Lasalocid
200	4	10000	300	200	1800	1000	30
	4	20000	2000	100	1900	1000	100
	6	6670	300	200	1800	667	30
	6	13340	1300	100	1900	667	65
500	10	10000	300	200	1800	1000	30
	10	20000	800	100	1900	1000	40

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.

**BLUE BIRD AUREO/BOV COCCI  
TYPE B Cattle Feed  
Medicated**

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle weighing up to 800 pounds.

**Active Drug Ingredients\***

Chlortetracycline ..... 4000 to 80000 g/ton\*\*  
Lasalocid ..... 181.8 to 3636.4 g/ton\*\*

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Mixing Directions**

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 500 to 4000 grams chlortetracycline per ton and 30 to 181.8 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
				CTC	Lasalocid
CTC	Lasalocid				
66667	3030	120	1880	4000	181.8
66667	3030	60	1940	2000	90.9
66667	3030	30	1970	1000	45.5
66667	3030	20	1980	667	30.3
40000	1818	200	1800	4000	181.8
40000	1818	100	1900	2000	90.9
40000	1818	50	1950	1000	45.5
40000	1818	33.3	1966.7	667	30.3

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

**NET WEIGHT ON BAG OR BULK**

=

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.

\*\* The ratio between chlortetracycline and lasalocid must be 22:1.