

**BLUE BIRD AUREO/BOV FE/ROG
TYPE B Confined Cattle Feed
Medicated**

For improved feed efficiency, increased rate of weight gain and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle fed in confinement for slaughter.

Active Drug Ingredients*

Chlortetracycline 4000 to 80000 g/ton
Lasalocid 100 to 80000 g/ton

Guaranteed Analysis

Crude protein, not less than %
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Mixing Directions

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 500 to 1200 grams chlortetracycline per ton and 25 to 30 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Body Wt. lb.	Feed Consumption lbs/head/day	Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
		CTC	Lasalocid			CTC	Lasalocid
800	20	8000	250	200	1800	800	25
	20	16000	600	100	1900	800	30
	24	6670	250	200	1800	667	25
	24	13340	600	100	1900	667	30
1000	20	10000	250	200	1800	1000	25
	20	20000	600	100	1900	1000	30
	25	8000	250	200	1800	800	25
	25	16000	560	100	1900	800	28

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.