

**BLUE BIRD AUREO/BOV ROG
TYPE B Pasture Cattle Feed
Medicated**

For increased rate of weight gain and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in pasture cattle (slaughter, stocker, feeder cattle, dairy and beef replacement heifers).

Active Drug Ingredients*

Chlortetracycline4000 to 80000 g/ton
Lasalocid400 to 80000 g/ton

Guaranteed Analysis

Crude protein, not less than%
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than%
Crude fiber, not more than%
Calcium, not less than%
Calcium, not more than%
Phosphorus, not less than%
Potassium, not less than%
Salt¹, not less than%
Salt¹, not more than%
Sodium², not less than%
Sodium², not more than%
Vitamin A¹, not less thanIU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Mixing Directions

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 500 to 4000 grams chlortetracycline per ton and 30 to 600 grams lasalocid per ton. The following table provides examples of mixing rates and amounts**:

Body Wt. lb.	Feed Consumption lbs/head/day	Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
		CTC	Lasalocid			CTC	Lasalocid
200	4	20000	2000	100	1900	1000	100
	6	6670	300	200	1800	667	30
	6	13340	1300	100	1900	667	65
	10	10000	300	200	1800	1000	30
500	10	20000	800	100	1900	1000	40

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.