

**BLUE BIRD AUREO/BOV FE  
TYPE C Confined Cattle Feed  
Medicated**

For improved feed efficiency and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle fed in confinement for slaughter.

**Active Drug Ingredients\***

Chlortetracycline ..... 500 to 2000 g/ton  
Lasalocid ..... 10 to 30 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Feed continuously in complete feed for not more than 5 days to provide 10 mg chlortetracycline per lb. body weight per day and not less than 100 mg nor more than 360 mg lasalocid per head per day. The following table provides examples of feeding rates and amounts:

Body Wt. lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/lb./day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
500	2	10	10	1000	100	20
	2	10	10	1000	150	30
	3	15	10	667	105	14
	3	15	10	667	225	30
1000	2	20	10	1000	100	10
	2	20	10	1000	300	30
	3	30	10	667	150	10
	3	30	10	667	360	24

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on this medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO/BOV FE/ROG  
TYPE C Confined Cattle Feed  
Medicated**

For improved feed efficiency, increased rate of weight gain and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle fed in confinement for slaughter.

**Active Drug Ingredients\***

Chlortetracycline ..... 500 to 1200 g/ton  
Lasalocid ..... 25 to 30 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Feed continuously in complete feed for not more than 5 days to provide 10 mg chlortetracycline per lb. body weight per day and not less than 250 mg nor more than 360 mg lasalocid per head per day. The following table provides examples of feeding rates and amounts:

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/lb./day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
800	2.5	20	10	800	250	25
	2.5	20	10	800	300	30
	3	24	10	667	300	25
	3	24	10	667	360	30
1000	2	20	10	1000	250	25
	2	20	10	1000	300	30
	2.5	25	10	800	312.5	25
	2.5	25	10	800	350	28

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on this medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO/BOV ROG  
TYPE C Pasture Cattle Feed  
Medicated**

For increased rate of weight gain and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in pasture cattle (slaughter, stocker, feeder cattle, dairy and beef replacement heifers).

**Active Drug Ingredients\***

Chlortetracycline ..... 500 to 4000 g/ton  
Lasalocid ..... 30 to 600 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Feed continuously on a hand-fed basis for not more than 5 days to provide 10 mg chlortetracycline per lb. body weight per day and not less than 60 mg or more than 300 mg lasalocid per head per day in at least 1 pound of feed for pasture cattle (slaughter, stocker, feeder cattle, dairy and beef replacement heifers). Daily intakes of lasalocid in excess of 200 mg/head/day have not been shown to be more effective than 200 mg/head/day. The following table provides examples of feeding rates and amounts:

Body Wt. lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/lb./day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
200	2	4	10	1000	60	30
	2	4	10	1000	200	100
	3	6	10	667	90	30
	3	6	10	667	195	65
500	2	10	10	1000	150	30
	2	10	10	1000	200	40

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on this medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO/BOV COCCI  
TYPE C Cattle Complete Feed  
Medicated**

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle weighing up to 800 pounds.

**Active Drug Ingredients\***

Chlortetracycline ..... 500 to 4000 g/ton\*\*  
Lasalocid ..... 30 to 181.8 g/ton\*\*

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Hand feed continuously for not more than 5 days to provide 10 mg chlortetracycline per lb. body weight per day and 1 mg lasalocid per 2.2 lb. body weight per day to cattle weighing up to 800 lb. with a maximum of 360 mg of lasalocid per head per day. The following table provides examples of feeding rates and amounts:

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/lb./day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
200	.5	1	10	4000	90.9	181.8
	1	2	10	2000	90.9	90.9
	2	4	10	1000	90.9	45.5
	3	6	10	667	90.9	30.3
800	.5	4	10	4000	360	180
	1	8	10	2000	360	90
	2	16	10	1000	360	45
	3	24	10	667	360	30

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on this medicated feed must bear a single concentration for each drug.

\*\* The ratio between chlortetracycline and lasalocid must be 22:1.