

**BLUE BIRD AUREO350 RESP/BOV FE
TYPE B Confined Cattle Feed
Medicated**

For improved feed efficiency and for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline in cattle fed in confinement for slaughter.

Active Drug Ingredients*

Chlortetracycline 100 to 80000 g/ton
Lasalocid 100 to 80000 g/ton

Guaranteed Analysis

Crude protein, not less than %
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Mixing Directions

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 25 to 100 grams chlortetracycline per ton and 10 to 30 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Body Wt. lb.	Feed Consumption lbs/head/day	Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
		CTC	Lasalocid			CTC	Lasalocid
	10	1400	600	100	1900	70	30
	15	467	140	200	1800	46.7	14
	15	933	600	100	1900	46.7	30
1000	20	350	100	200	1800	35	10
	20	700	600	100	1900	35	30
	28	250	100	200	1800	25	10
	28	500	500	100	1900	25	25

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.