

**BLUE BIRD AUREO350 RESP/BOV COCCI
TYPE B Cattle Feed
Medicated**

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline in beef cattle weighing up to 800 pounds.

Active Drug Ingredients*

Chlortetracycline2800 to 80000 g/ton
Lasalocid 181.8 to 3636.4 g/ton

Guaranteed Analysis

Crude protein, not less than %
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Mixing Directions

Mix the Type B medicated feed with non-medicated ingredients to manufacture one ton of complete cattle feed containing 25 to 2800 grams chlortetracycline per ton and 30 to 181.8 grams lasalocid per ton. The following table provides examples of mixing rates and amounts:

Type B Concentration, g/ton		Type B per ton of Type C lbs	Lbs Non-medicated feed per ton of Type C feed	Type C Concentration, g/ton	
				CTC	Lasalocid
CTC	Lasalocid				
11666.67	3030	120	1880	700	181.8
11666.67	3030	60	1940	350	90.9
11666.67	3030	30	1970	175	45.5
11666.67	3030	20	1980	116.7	30.3
7000	1818	200	1800	700	181.8
7000	1818	100	1900	350	90.9
7000	1818	50	1950	175	45.5
7000	1818	33.3	1966.7	116.7	30.3

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle. Do not allow horses or other equines access to Type B feeds containing lasalocid as ingestion may be fatal.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on formulated Type B Medicated Feed must bear a single concentration for each drug.