

**BLUE BIRD AUREO350 RESP/BOV FE  
TYPE C Confined Cattle Feed  
Medicated**

For improved feed efficiency and for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline in cattle fed in confinement for slaughter.

**Active Drug Ingredients\***

Chlortetracycline ..... 25 to 100 g/ton  
Lasalocid ..... 10 to 30 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Feed continuously in complete feed at a rate of 350 mg chlortetracycline and not less than 100 mg nor more than 360 mg lasalocid per head daily. The following table provides examples of feeding rates and amounts:

Body Wt. lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/head/day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
500	1.4	7	350	100	105	30
	2	10	350	70	100	20
	3	15	350	46.7	105	14
	3	15	350	46.7	225	30
1000	2	20	350	35	100	10
	2	20	350	35	300	30
	2.8	28	350	25	140	10
	2.8	28	350	25	350	25

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on this medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO350 RESP/BOV FE/ROG  
TYPE C Confined Cattle Feed  
Medicated**

For improved feed efficiency, increased rate of weight gain and for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline in cattle fed in confinement for slaughter.

**Active Drug Ingredients\***

Chlortetracycline ..... 25 to 42.2 g/ton  
Lasalocid ..... 25 to 30 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Feed continuously in complete feed at a rate of 350 mg chlortetracycline and not less than 250 mg nor more than 360 mg lasalocid per head daily. The following table provides examples of feeding rates and amounts:

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/head/day	Feeding Rate, g/ton	Amount mg/head/day	Feeding Rate g/ton
800	2.1	16.6	350	42.2	250	30
	2.5	20	350	35	250	25
	2.5	20	350	35	300	30
	3	24	350	29.2	360	30
1000	2	20	350	35	250	25
	2	20	350	35	300	30
	2.5	25	350	28	312.5	25
	2.5	25	350	28	350	28

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO350 RESP/BOV ROG  
TYPE C Pasture Cattle Feed  
Medicated**

For increased rate of weight gain and for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline in pasture cattle (slaughter, stocker, feeder cattle, beef replacement heifers).

**Active Drug Ingredients\***

Chlortetracycline ..... 25 to 700 g/ton  
Lasalocid ..... 30 to 600 g/ton

**Guaranteed Analysis**

Crude protein, not less than ..... %  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %  
Vitamin A<sup>1</sup>, not less than ..... IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Feed continuously on a hand-fed basis at a rate of 350 mg chlortetracycline and not less than 60 mg or more than 300 mg lasalocid per head daily in at least 1 pound of feed for pasture cattle (slaughter, stocker, feeder cattle, dairy and beef replacement heifers). Daily lasalocid intakes in excess of 200 mg/head/day in pasture cattle have not been shown to be more effective than 200 mg lasalocid/head/day. The following table provides examples of feeding rates and amounts:

Body Wt. lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/head/day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
200	0.5	1	350	700	300	600
	2	4	350	175	200	100
	3	6	350	116.7	90	30
	3	6	350	116.7	195	65
500	2	10	350	70	150	30
	2	10	350	70	200	40

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on this medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO350 RESP/BOV COCCI  
TYPE C Cattle Complete Feed  
Medicated**

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline in beef cattle weighing up to 800 pounds.

**Active Drug Ingredients\***

Chlortetracycline .....25 to 2800 g/ton  
Lasalocid .....30 to 181.8 g/ton

**Guaranteed Analysis**

Crude protein, not less than .....%  
(This includes not more than \_\_\_% equivalent crude protein from non-protein nitrogen)<sup>1</sup>  
Crude fat, not less than .....%  
Crude fiber, not more than .....%  
Calcium, not less than .....%  
Calcium, not more than .....%  
Phosphorus, not less than.....%  
Potassium, not less than .....%  
Salt<sup>1</sup>, not less than .....%  
Salt<sup>1</sup>, not more than.....%  
Sodium<sup>2</sup>, not less than .....%  
Sodium<sup>2</sup>, not more than .....%  
Vitamin A<sup>1</sup>, not less than .....IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**Feeding Directions**

Hand feed continuously at a rate of 350 mg chlortetracycline per head and 1 mg lasalocid per 2.2 lb. body weight daily to cattle with a maximum of 360 mg of lasalocid per head per day. The following table provides examples of feeding rates and amounts:

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb./head/day	Amount mg/head/day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
200	0.5	1	350	700	90.9	181.8
	1	2	350	350	90.9	90.9
	2	4	350	175	90.9	45.5
500	2	10	350	70	227.3	45.5
	2.5	12.5	350	56	227.3	36.4
800	2	16	350	43.8	360	45
	3	24	350	29.2	360	30

**Warning**

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

**NET WEIGHT ON BAG OR BULK**

Blue Bird Feed Mill  
Robin, IN 12345

\*Final printed label on this medicated feed must bear a single concentration for each drug.