

**BLUE BIRD AUREO350 ANA/BOV FE
TYPE C Confined Cattle Feed
Medicated**

For improved feed efficiency and for control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline in cattle under 700 pounds fed in confinement for slaughter.

Active Drug Ingredients*

Chlortetracycline 25 to 100 g/ton
Lasalocid 10 to 30 g/ton

Guaranteed Analysis

Crude protein, not less than %
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Feeding Directions

Feed continuously in complete feed at a rate of 350 mg chlortetracycline and not less than 100 mg nor more than 360 mg lasalocid per head daily. The following table provides examples of feeding rates and amounts:

| Body Wt. lb. | Feed Consumption | | Chlortetracycline | | Lasalocid | |
|-----------------|------------------|--------------|-----------------------|-----------------------|-----------------------|--------------------------|
| | % Body Wt. | lb./head/day | Amount mg/head/day | Feeding Rate g/ton | Amount mg/head/day | Feeding Rate g/ton |
| 500 | 1.4 | 7 | 350 | 100 | 105 | 30 |
| | 2 | 10 | 350 | 70 | 150 | 30 |
| | 3 | 15 | 350 | 46.7 | 105 | 14 |
| | 3 | 15 | 350 | 46.7 | 225 | 30 |
| 690 | 2.9 | 20 | 350 | 35 | 100 | 10 |
| | 2.9 | 20 | 350 | 35 | 300 | 30 |

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on this medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO350 ANA/BOV FE/ROG
TYPE C Confined Cattle Feed
Medicated**

For improved feed efficiency, increased rate of weight gain and for control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline in cattle under 700 pounds fed in confinement for slaughter.

Active Drug Ingredients*

Chlortetracycline 25 to 42.2 g/ton
Lasalocid 25 to 30 g/ton

Guaranteed Analysis

Crude protein, not less than %
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Feeding Directions

Feed continuously in complete feed at a rate of 350 mg chlortetracycline and not less than 250 mg nor more than 360 mg lasalocid per head daily. The following table provides examples of feeding rates and amounts:

| Body Wt. Lb. | Feed Consumption | | Chlortetracycline | | Lasalocid | |
|-----------------|------------------|--------------|-----------------------|-----------------------|-----------------------|--------------------------|
| | % Body Wt. | lb./head/day | Amount mg/head/day | Feeding Rate g/ton | Amount mg/head/day | Feeding Rate g/ton |
| 675 | 2.5 | 17 | 350 | 41.2 | 251 | 29.5 |
| | 3.1 | 21 | 350 | 33.33 | 263 | 25 |
| | 3.1 | 21 | 350 | 33.33 | 315 | 30 |

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on formulated Type C Medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO350 ANA/BOV ROG
TYPE C Pasture Cattle Feed
Medicated**

For increased rate of weight gain and for control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline in pasture cattle under 700 pounds (slaughter, stocker, feeder cattle, beef replacement heifers).

Active Drug Ingredients*

Chlortetracycline 25 to 700 g/ton
Lasalocid 30 to 600 g/ton

Guaranteed Analysis

Crude protein, not less than %
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Feeding Directions

Feed continuously on a hand-fed basis at a rate of 350 mg chlortetracycline and not less than 60 mg or more than 300 mg lasalocid per head daily in at least 1 pound of feed for pasture cattle (slaughter, stocker, feeder cattle, dairy and beef replacement heifers). Daily lasalocid intakes in excess of 200 mg/head/day in pasture cattle have not been shown to be more effective than 200 mg lasalocid/head/day. The following table provides examples of feeding rates and amounts:

| Body Wt. lb. | Feed Consumption | | Chlortetracycline | | Lasalocid | |
|-----------------|------------------|--------------|-----------------------|-----------------------|-----------------------|--------------------------|
| | % Body Wt. | lb./head/day | Amount Mg/head/day | Feeding Rate g/ton | Amount mg/head/day | Feeding Rate g/ton |
| 200 | 0.5 | 1 | 350 | 700 | 300 | 600 |
| | 2 | 4 | 350 | 175 | 200 | 100 |
| | 3 | 6 | 350 | 116.7 | 90 | 30 |
| | 3 | 6 | 350 | 116.7 | 195 | 65 |
| 500 | 2 | 10 | 350 | 70 | 150 | 30 |
| | 2 | 10 | 350 | 70 | 200 | 40 |

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on this medicated feed must bear a single concentration for each drug.

**BLUE BIRD AUREO350 ANA/BOV COCCI
TYPE C Cattle Complete Feed
Medicated**

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline in beef cattle under 700 pounds.

Active Drug Ingredients*

Chlortetracycline25 to 2800 g/ton
Lasalocid30 to 181.8 g/ton

Guaranteed Analysis

Crude protein, not less than %
(This includes not more than ___% equivalent crude protein from non-protein nitrogen)¹
Crude fat, not less than %
Crude fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Potassium, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %
Vitamin A¹, not less than IU/lb

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

Feeding Directions

Hand feed continuously at a rate of 350 mg chlortetracycline per head and 1 mg lasalocid per 2.2 lb. body weight daily to cattle with a maximum of 360 mg of lasalocid per head per day. The following table provides examples of feeding rates and amounts:

| Body Wt. Lb. | Feed Consumption | | Chlortetracycline | | Lasalocid | |
|-----------------|------------------|--------------|-----------------------|-----------------------|-----------------------|-----------------------|
| | % Body Wt. | lb./head/day | Amount mg/head/day | Feeding Rate g/ton | Amount mg/head/day | Feeding Rate g/ton |
| 200 | 0.5 | 1 | 350 | 700 | 90.9 | 181.8 |
| | 1 | 2 | 350 | 350 | 90.9 | 90.9 |
| | 2 | 4 | 350 | 175 | 90.9 | 45.5 |
| 500 | 2 | 10 | 350 | 70 | 227.3 | 45.5 |
| | 2.5 | 12.5 | 350 | 56 | 227.3 | 36.4 |
| 700 | 2 | 14 | 350 | 50 | 318.2 | 45.5 |
| | 3 | 21 | 350 | 33.3 | 318.2 | 30.3 |

Warning

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345

*Final printed label on this medicated feed must bear a single concentration for each drug.