

Now Available In The U.S.!



Ready To Use

- ✓ Simply dip the full length of the teats after taking off the milking unit – *it's that easy.*
- ✓ Do not dilute or mix with other chemicals.
- ✓ Discard if contamination occurs.
- ✓ Always use a clean dip cup.



Available in 5-gallon and 55-gallon sizes.

KENOSTART®



KENOSTART®

ALPHARMA
Animal Health

Alpharma Inc.
One Executive Drive
Fort Lee, NJ 07024, USA

tel: (800) 821-5811
www.alpharma.com

References

1. Determination of basic bactericidal activity according to standard EN 1040, chemical disinfectants and antiseptics Phase 1. March 23-April 9, 2004. Belgium.
2. Determination of bactericidal activity according to standard EN 1656; chemical disinfectants and antiseptics used in veterinary field Phase 2, step1. March 25-April 9, 2004. Belgium.
3. Teat conditioning trial report for an experimental post - milking teat dip. November, 2003. Belgium.

KENOSTART is a registered trademark of CID Lines.
Manufactured under ISO 9001:2000 and GMP Pharma registration by
CID Lines n.v., Belgium - www.cidlines.com.
©2006 Alpharma Inc. All rights reserved. CD0519

Delivering Total Security Around The Teat

The Premium Iodine-Based Teat Dip That Delivers Advanced Protection As It Conditions



Not Just Another Iodine Dip

- Advanced protection against environmental- and mastitis-causing bacteria
- Specially formulated – higher level of active stable iodine and a specific surfactant for cleaning without irritation
- Protective barrier significantly heals and conditions – keeping skin supple and soft
- Non-drip for greater efficiency – uniform film improves viscosity

World-Class Teat Protection


- **BASIC BACTERICIDAL ACTIVITY** effective against *Pseudomonas aeruginosa* and *Staphylococcus aureus*¹
- **BACTERICIDAL ACTIVITY IN THE VETERINARY FIELD CONDITIONS** effective against *P. aeruginosa*, *S. aureus*, *Proteus vulgaris*, and *Enterococcus hirae* in contact times of 5 and 30 minutes with milk as the interfering substance²
- **SPECIFIC MASTITIS BACTERIA** effective against *Escherichia coli*, *Streptococcus uberis*, *Streptococcus agalactiae*, and *Streptococcus dysgalactiae* in contact times of 5 and 30 minutes with milk as the interfering substance²



Teat Conditioning – Field-Proven³

- 8-week study
- KENOSTART 2 times per day after milking improved average teat score from 3 to less than 2
- At end of trial teats were smooth, free from scales, cracks, or chapping
- Teat end sphincter was smooth with no evidence of irritation

Teat Condition Scoring: KENOSTART³



4. Teat skin is chapped and cracked. Redness, indicating inflammation, is present. Numerous warts may be present.



3. Teat skin is chapped. Some small warts may be present.



2. Teat skin shows some evidence of scaling.



1. Teat skin is smooth, free from scales, cracks, or chapping.

AOAC Testing Confirms Antibacterial Activity Against These Strains:^{3*}

<i>Pseudomonas aeruginosa</i>	<i>Klebsiella pneumoniae</i>
<i>Staphylococcus aureus</i>	<i>Streptococcus bovis</i>
<i>Proteus vulgaris</i>	<i>Corynebacterium bovis</i>
<i>Escherichia coli</i>	<i>Streptococcus agalactiae</i>
<i>Citrobacter freundii</i>	<i>Streptococcus dysgalactiae</i>
<i>Enterobacter aerogenes</i>	<i>Streptococcus uberis</i>

* At a concentration of 94% (v/v) at 30°C with 30 sec. contact time and under 5% whole milk conditions.

KENOSTART[®]

Delivering Total Security Around The Teat