

Sul-Q-Nox™

31.92%
Sulfaquinoxaline

Drinking water
administration

- Concentrated liquid is convenient for water medication
- Effective sulfa treatment for major poultry diseases
- Suitable for automatic water proportioner systems or tanks
- Convenient sizes for either tank or proportioner administration

Composition:

Liquid Sul-Q-Nox contains 31.92% sulfaquinoxaline as sodium and potassium salts.

Indications/Dose:

- **Liquid Sul-Q-Nox** is to be used as an aid in the control of coccidiosis outbreaks in chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima* and *E. brunetti* susceptible to sulfaquinoxaline.
- For turkeys, it is to be used as an aid in the control of outbreaks of coccidiosis caused by *Eimeria meleagrimitis* and *E. adenoides* susceptible to sulfaquinoxaline.
- In chickens and turkeys, it aids in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline and fowl typhoid caused by *Salmonella gallinarum* susceptible to sulfaquinoxaline.
- In cattle and calves, use for the control and treatment of outbreaks of coccidiosis caused by *Eimeria bovis* or *E. zurnii*.

For more information on putting Liquid Sul-Q-Nox to work for you, contact your nearest Alpha representative.

Alpha – The performance enhancement people

Sul-Q-Nox™



Alpha Inc.
One Executive Drive
Fort Lee, NJ 07024, USA

Tel (201) 947-7774
Fax (201) 585-0429
<http://www.alpha.com>

Sul-Q-Nox is a trademark of Alpha Inc.

Copyright ©2001 by Alpha Inc.

Proportioner Mixing Instructions:

Add the following amount of Liquid Sul-Q-Nox to water to make a 1 gallon stock solution for proportioner set to meter at the rate of one ounce per gallon (1:128).

Species	Disease	Amount per Gallons of Stock Solution	Treatment	Withdrawal
Turkeys ¹	Coccidiosis	13 fl. oz. (0.025%)	Give for 2 days, skip 3 days, give for 2 days, skip 3 days, give for 2 more days	10 Days
	Fowl Cholera	20.5 fl. oz. (0.04%)	2-3 days, repeat if disease recurs	10 Days
	Fowl Typhoid	20.5 fl. oz. (0.04%)	2-3 days, repeat if disease recurs	10 Days
Chickens ¹	Coccidiosis	20.5 fl. oz. (0.04%) Then 13 fl. oz. (0.025%) 13 fl. oz. (0.025%)	Give for 2-3 days, skip 3 days Then Give 2 more days Give 2 more days	10 Days 10 Days 10 Days
	<i>If bloody droppings appear:</i>			
	Fowl Cholera	20.5 fl. oz. (0.04%)	2-3 days, repeat if disease recurs	10 Days
	Fowl Typhoid	20.5 fl. oz. (0.04%)	2-3 days, repeat if disease recurs	10 Days
Cattle & Calves ²	Coccidiosis	7.75 fl. oz. (0.015%)	3-5 days	10 Days (not established in pre-ruminating calves)

Tank Mixing Instructions:

Add the appropriate amount of product to a tank of adequate volume and then add the amount of water listed below to make treatment solutions. See above for indications and use levels.

Solution Levels	Amount of Product		
	1 Ounce	50 Ounce	1 Gallon
0.04%	6.2 gallons water	311 gallons water	797 gallons water
0.025%	10.0 gallons water	498 gallons water	1276 gallons water
0.015%	16.6 gallons water	831 gallons water	2127 gallons water

¹ Do not medicate chickens or turkeys producing eggs for human consumption.

² Do not use in calves to be processed for veal. Do not use in lactating dairy cattle.

See package label for complete use directions and warnings.

Package size:

4 x 1 gallon

6 x 50 ounces